

given to charge Deposit Account No. 23-1701 in the amount of Four Hundred Dollars (\$400.00)

to cover the corresponding extension fee pursuant to 37 C.F.R. §§1.17(a)(2) and 1.136(a).

OFFICIAL**FAX RECEIVED**

AUG 16 2002

GROUP 1600**IN THE CLAIMS:**


Substitute claims 1, 18, 26 and 27 with amended claims 1, 18, 26 and 27:

81
E1
1. (Four times amended) A method of treatment for improving the inhibition of gastric acid secretion which comprises administering to a host in need thereof an oral pharmaceutical formulation comprising a therapeutically effective amount of an acid labile H^+ , K^+ -ATPase inhibitor, wherein the method induces an extended blood plasma profile of the H^+ , K^+ -ATPase inhibitor, and the H^+ , K^+ -ATPase inhibitor is a compound of the formula I




wherein

Het₁ is




or




Het₂ is


*Sub
J1
cont*




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

*E1
concl.* R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxycarbonyl, oxazolyl, trifluoroalkyl, or adjacent groups R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and


R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl.

18. (Thrice amended) A method of treatment for improving the inhibition of gastric acid secretion which comprises administering to a host in need thereof an oral pharmaceutical formulation comprising a therapeutically effective amount of an acid labile H^+ , K^+ -ATPase inhibitor, wherein the method induces an extended blood plasma profile of the H^+ , K^+ -ATPase inhibitor, and the H^+ , K^+ -ATPase inhibitor is a compound of the formula I




wherein

Het₁ is




or




Het₂ is


*Sub
32
cont*




or



X =



or



*Ed
Concl* wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxy carbonyl, oxazolyl, trifluoroalkyl, or adjacent groups R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and

R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl


with the proviso that the H⁺, K⁺-ATPase inhibitor is not pantoprazole.

26. (Twice amended) A method for improving the treatment of gastrointestinal disorders associated with excess acid secretion which comprises administering to a host in need thereof an oral pharmaceutical formulation comprising a therapeutically effective amount of an acid labile H^+ , K^+ -ATPase inhibitor, wherein the method induces an extended blood plasma profile of the H^+ , K^+ -ATPase inhibitor, and the H^+ , K^+ -ATPase inhibitor is a compound of the formula I




wherein


Het_1 is




or




Het_2 is




or



Sub 33 cont
X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;


R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxycarbonyl, oxazolyl, trifluoroalkyl, or adjacent groups R₆-R₉ form ring structures which may be further substituted;

93 R₁₀ is hydrogen or forms an alkylene chain together with R₃; and

R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl.


27. (Twice amended) A method for improving the treatment of gastrointestinal disorders associated with excess acid secretion which comprises administering to host in need thereof an oral pharmaceutical formulation comprising a therapeutically effective amount of an acid labile H⁺, K⁺-ATPase inhibitor, wherein the method induces an extended blood plasma profile of the H⁺, K⁺-ATPase inhibitor, and the H⁺, K⁺-ATPase inhibitor is a compound of the formula I




I

wherein


Het₁ is




or




Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;